

University of Montana

## ScholarWorks at University of Montana

---

University of Montana Conference on Undergraduate Research (UMCUR)

---

Apr 15th, 3:00 PM - 4:00 PM

### Photoionization Cross-Sections and Rydberg Resonance Identifications of Br<sup>+</sup>

Dylan Gross

*Department of Physics and Astronomy, University of Montana, [dylan.gross@umconnect.umt.edu](mailto:dylan.gross@umconnect.umt.edu)*

Follow this and additional works at: <https://scholarworks.umt.edu/umcur>

## Let us know how access to this document benefits you.

---

Gross, Dylan, "Photoionization Cross-Sections and Rydberg Resonance Identifications of Br<sup>+</sup>" (2016).  
*University of Montana Conference on Undergraduate Research (UMCUR)*. 18.  
<https://scholarworks.umt.edu/umcur/2016/pm posters/18>

This Poster is brought to you for free and open access by ScholarWorks at University of Montana. It has been accepted for inclusion in University of Montana Conference on Undergraduate Research (UMCUR) by an authorized administrator of ScholarWorks at University of Montana. For more information, please contact [scholarworks@mso.umt.edu](mailto:scholarworks@mso.umt.edu).



# Photoionization Cross-Sections and Rydberg Resonance Identifications of $\text{Br}^+$

Dylan Gross<sup>1</sup>, D. A. Macaluso<sup>1</sup>, A. Aguilar<sup>2</sup>, R. C. Bilodeau<sup>3</sup>, A. L. D. Kilcoyne<sup>2</sup>, and N. C. Sterling<sup>4</sup>

<sup>1</sup>Department of Physics and Astronomy, University of Montana, Missoula, MT.

<sup>2</sup>The Advanced Light Source, Lawrence Berkeley National Laboratory, Berkeley, CA. <sup>3</sup>Department of Physics, University of Connecticut, Storrs, CT.

<sup>4</sup>Department of Physics, University of West Georgia, Carrollton, GA.



## MOTIVATION

Astrophysicists often need atomic data for elements that are detected in astrophysical nebulae. Because trans-iron elements have recently been detected, specifically in planetary nebulae, and have very little atomic data, our goal has been to determine absolute single photoionization for trans-iron elements. The present analysis' purpose is to find photoionization cross-sections, to find and validate direct ionization thresholds and excited state energy levels and to identify Rydberg series in photo-ion spectroscopy of  $\text{Br}^+$  ions.

A few examples of planetary nebula can be seen to the right.




NGC 5189 Nebula




Red Spider Nebula (NGX 6537)

## EXPERIMENT

The Advanced Light Source at Lawrence Berkeley National Laboratory uses a linear electron accelerator and booster ring placed within the main storage ring.




**Experimental Technique:** Positively charged Br ions are created in the **electron-cyclotron-resonance (ECR) ion source**.  $\text{Br}^+$  ions are mass-charged selected using a **60° analyzing magnet** and then merged with the **monochromatized photon beam**. **3 translating slit scanners** within the **29.4 cm long biased interaction region** are used to tune the two beams to have a good overlap. A **45° demerging magnet** then separates the  $\text{Br}^{2+}$  photo ions from the primary ion beam and are sent to the **product ion detector** [1],[2],[3],[4].



Beamline 10.0.1.2 End Station

## RESULTS




Absolute single photoionization cross section measurements are represented by open circles with  $\pm 20\%$  error bars. These exact values are also presented in table 1.


Table 1.

Energy (eV)	Cross Section (Mb)	Uncertainty ( $\pm$ )
18.141	13.9116	2.78232
22.386	17.5	3.5
23.69	13.7	2.74
26	8.47	1.694
28	637	1.274
30	5.03	1.006
32	3.7	0.74

Because these absolute cross section measurements had not been studied to this degree of precision, these data will help astrophysicists to determine abundances of  $\text{Br}^+$  ions in planetary nebulae.



**Panel A** shows the one identified Rydberg series which originates from the  $3P_{3/2}$  parent ion state of  $\text{Br}^+$  and converges to the  $2P_{3/2}$  final product ion state of  $\text{Br}^{2+}$ . Ionization thresholds and additional Rydberg series are still being determined. **Panel B** shows an example of what our  $\text{Br}^+$  data should look like after further analysis of the spectrum.



Dylan Gross would like to thank and acknowledge MSGC for funding his current data analysis research and Dr. Macaluso for facilitating the research.

### References

- [1] N. C. Sterling, D. A. Esteves, R. C. Bilodeau, A. L. D. Kilcoyne, E. C. Red, R. A. Phaneuf, and A. Aguilar, J. Phys. B 44, 025701 (2011). [2] D. A. Esteves, R. C. Bilodeau, N. C. Sterling, R. A. Phaneuf, A. L. D. Kilcoyne, E. C. Red, and A. Aguilar, Phys. Rev. A 92, 063424 (2015). [3] D. A. Esteves, A. Aguilar, A. L. D. Kilcoyne, E. C. Red, R. C. Bilodeau, R. A. Phaneuf, N. C. Sterling, and B. M. McLaughlin, Phys. Rev. A 92, 063424 (2015). [4] D. A. Esteves, R. C. Bilodeau, R. A. Phaneuf, A. L. D. Kilcoyne, E. C. Red, and A. Aguilar, J. Phys. B 45, 115201 (2012). [5] <http://scitechdaily.com/hubble-views-planetary-nebula-ngc-5189/> (2012). [6] <http://annesastronomynews.com/photo-gallery-ii/nebulae-clouds/the-red-spider-nebula-ngc-6537/> [7] <https://www.ickr.com/photos/advancedlightsource/4206760015/in/album-72157623046767608/> (2005).